


Effect of Water Deficit Stress on Reproductive Stage of Durum Wheat Near Isogenic Lines Carrying the *NAM-B1* Gene



Farah O. AL Mbideen¹, Abeer Aburumman², Rabea Al-Sayaydeh³, Randa N. Albdaiwi⁴ and Ayed M. Al-Abdallat^{1,*} 

¹Department of Horticulture and Crop Science, School of Agriculture, The University of Jordan, Amman 11942, Jordan

²Department of Plant Biotechnology, National Agricultural Research Center - NARC, Amman, Jordan

³Department of Plant Production, Shoubak Collage, Al-Balqa Applied University, Al-Salt 19117, Jordan

⁴Department of Allied Medical Sciences, Zarqa University College, Al-Balqa Applied University, Al-Salt 19117, Jordan

Abstract:

Background: In wheat, the NAC family plays a crucial role in conferring cellular responses to different abiotic stresses, particularly drought. The functional *NAM-B1* allele in durum wheat accelerates senescence, while its homologous genes contribute to improved water deficit stress tolerance. Therefore, the main objective of this study was to investigate the role of the *NAM-B1* gene in durum wheat subjected to water deficit stress at two reproductive stages.

Materials and Methods: Three Near-Isogenic Lines carrying a functional *NAM-B1* allele and their recurrent parent, Langdon (LDN), were used in this study. The study assessed NILs and LDN performance under varied water conditions (well-watered, 60%, and 80% water deficit of pot capacity) during flag leaf and anthesis stages. The agronomic performance of the three NILs and LDN was assessed for 16 traits, including plant height, grain yield, and thousand kernel weight. Additionally, physiological measurements, including stomatal resistance, chlorophyll content (SPAD values), and chlorophyll fluorescence (*Fv/Fm*) were taken.

Results: Significant genotypic effects were observed on seven agronomic traits, while 15 traits were influenced by the water stress treatment. The NILs exhibited accelerated maturity and a shorter grain-filling period, which were particularly pronounced under stress conditions. Severe water deficit resulted in reduced grain weight and thousand-grain weight in tested genotypes. Interestingly NILs carrying functional *NAM-B1* showed taller plants and had higher tiller and spike numbers when compared to LND. Significant genotypic effects were observed for seven traits, and water stress treatments affected 15 traits. The NILs exhibited accelerated maturity and reduced grain weight under severe water deficit. Physiological measurements showed genotype and water-deficit differences, with NIL #504 displaying higher SPAD values, particularly under stress conditions, and a significant genotype X treatment effect was observed for stomatal resistance at the anthesis stage.

Discussion: Limited treatment × genotype interactions for most traits highlight the complexity of drought tolerance in the tested lines, which might be attributed to its quantitative nature influenced by multiple genes. The study emphasizes the necessity for future research to explore the role of other *NAM*-related genes in response to water deficit stress and their interactive effects with the *NAM-B1* gene at different growth stages in wheat plants subjected to diverse stress conditions.

Conclusion: In conclusion, this study reveals pronounced genotypic effects, particularly in agronomic traits, such as maturation rate and grain filling period, under severe water deficit stress. Future studies are needed to understand the genetic mechanisms mediated by *NAM-B1* genes in accelerating maturity in response to stress.

Keywords: Drought, Durum wheat, Senescence, Stomatal resistance, Transcription factors, Stress.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: March 03, 2024
Revised: April 19, 2024
Accepted: April 26, 2024
Published: June 05, 2024

*Address correspondence to this author at the Department of Horticulture and Crop Science, School of Agriculture, The University of Jordan, Amman 11942, Jordan; E-mail: a.alabdallat@ju.edu.jo

Cite as: Mbideen F, Aburumman A, Al-Sayaydeh R, Albdaiwi R, Al-Abdallat A. Effect of Water Deficit Stress on Reproductive Stage of Durum Wheat Near Isogenic Lines Carrying the *NAM-B1* Gene. Open Agric J, 2024; 18: e18743315314060. <http://dx.doi.org/10.2174/0118743315314060240516065334>



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Wheat (*Triticum* spp.) is one of the most important cultivated crops in the world. It ranks first among cultivated plants in terms of cultivated areas around the globe [1]. It is considered a major staple food for over 35% of the world's population, providing more than 20% of their daily protein and calorie needs [2]. Durum wheat (*Triticum turgidum* ssp. *durum*) constitutes approximately 10% of the total cultivated wheat area and is predominantly cultivated in the Mediterranean region under rainfed conditions [3]. Its distinctive vitreous grain structure and high protein content make it suitable for various uses, including the production of pasta, semolina, flatbreads, couscous, and bulgur [4, 5].

Climate change and associated conditions, including changes in temperature, precipitation patterns, and the frequency and intensity of extreme weather events [6], are expected to have a significant impact on the productivity of many crops, including wheat [7]. Drought is one of the most common environmental stresses that can have a significant impact on the growth and development of wheat plants [8]. Under such conditions, wheat production and quality are adversely affected, especially during critical growth stages, such as heading and grain-development phases [9]. For instance, severe drought stress in wheat plants during the grain-filling stage reduced yields by 44% [10]. In another study, prolonged mild drought stress at the heading stage reduced grain yield by 58% [11]. Several yield-related traits, including spike length, grains per spike, total biomass, and grain yield, are expected to be affected under drought conditions [6, 7]. Hence, there is an urgent need to breed new wheat varieties with climate-smart traits capable of tolerating various abiotic stresses, including drought [12].

The identification and utilization of key regulatory genes associated with stress tolerance is considered a priority for producing new varieties with improved drought tolerance [13]. In this regard, transcription factors play essential roles in controlling multiple gene expression pathways and are considered key regulatory genes for many cellular processes [14]. In plants, the NAC (NAM-ATAF-CUC2) or NAM (No Apical Meristem) transcription factor family plays a major role in regulating various cellular processes, such as hormonal signaling, plant development, and responses to biotic and abiotic stresses [15]. Additionally, *NAM* genes play important

roles in regulating gene expression changes during leaf senescence, and several *NAM* genes have been found to promote Reactive oxygen species (ROS) accumulation and significantly accelerate age-dependent as well as drought- and Abscisic acid (ABA)-induced leaf senescence [16, 17]. *NAM* genes associated with abiotic stress tolerance have been cloned and successfully used to improve stress tolerance, particularly drought tolerance, in different crop species [18-21].

In wheat, the *NAM-B1* gene was cloned as a major locus for grain protein content and senescence [22]. The *NAM-B1* gene is considered a positive regulator of senescence and is associated with increased protein, Fe, Mn, and Zn contents in wheat grains [23]. Kaznina *et al.* [24] demonstrated that wheat plants carrying a functional allele of the *NAM-B1* gene exhibit good growth under zinc deficiency conditions without any decrease in yield. In nature, this gene is only functional in wild emmer and a few domesticated emmer wheat varieties, while frame-shift mutations or deletions in its coding sequence have been identified in modern bread or durum wheat genotypes, resulting in loss-of-function alleles and delayed senescence [21, 25]. This gene has been introgressed into new durum and bread wheat cultivars, and near-isogenic lines (NILs) carrying *NAM-B1* have been developed [21, 26, 27]. Furthermore, lines carrying a functional copy of *NAM-B1* have shown higher yield potential compared to local checks across different environments, including dry areas [28]. This could be attributed to the early maturity and abiotic stress tolerance associated with *NAM-B1* introgression.

Recent phylogenetic analysis of the NAC family in wheat identified several stress-responsive gene candidates, including two *NAM-B1* homologous genes (*NAM-A1* and *NAM-D1*) [29, 30]. The overexpression of *TaSNAC8-6A* (*NAM-A1*) in wheat plants has resulted in improved drought tolerance and enhanced water use efficiency, thus confirming its major role in stress tolerance [29]. Additionally, a natural allele called *TaSNAC8-6AIn-313*, which was found to be highly expressed under stress conditions, has been found to confer drought tolerance at the seedling stage in selected bread wheat genotypes [29]. These findings highlight the potential role of *NAM-A1* and its homologous genes in conferring drought tolerance in wheat plants. This study aimed to investigate the role of *NAM-B1*, a homologous gene to the stress-responsive *NAM-A1* gene, in conferring

tolerance to water deficit stress at two reproductive stages by using NILs carrying the *NAM-B1* allele. For this purpose, we evaluated the agronomic performance of the selected NILs and analyzed their physiological responses under water deficit stress conditions. The results obtained from this study would contribute to ongoing research efforts aimed at dissecting the role of *NAM-B1* in enhancing drought tolerance in wheat and improving crop productivity in water-limited environments.

2. MATERIALS AND METHODS

2.1. Plant Material

In this study, three durum wheat NILs, namely, IR51-8 (PI 656794), IR17-47 (PI 656795), and 504 (PI 656796) were obtained from the U.S. Department of Agriculture - Agricultural Research Service (USDA-ARS) national small grains collection. The three lines were developed at UC Davis (CA, USA). They were selected from a cross between the durum wheat cultivar Langdon (LDN) and RSL65, a recombinant inbred chromosomal line of Langdon carrying the *NAM-B1* gene, which was used in this study [21, 26]. The selected NILs carry a chromosome 6BS segment with a functional *NAM-B1* gene from *T. turgidum* subsp. *dicoccoides* (accession FA15-3), while the rest of their genome resembles LDN. The durum wheat cultivar LDN (Citr 13165) was used as a control. It carries a nonfunctional *NAM-B1* gene resulting from a 1 bp indel at the start of the gene, causing a frame-shift mutation and a non-functional allele [21].

2.2. Water Deficit Experiment

The water deficit stress experiment was conducted in the greenhouse facility at the School of Agriculture, The University of Jordan, starting from November 1st, 2019, until the plants reached physiological maturity (Zadoks scale: GS87) [30, 31]. Two seeds from each genotype were sown in a 6 L plastic pot filled with 5.5 kg of acid-washed sand. The pots were initially watered until they reached their “pot capacity,” estimated by adding 500 ml of distilled water to reach a pot weight of 6 kg. Subsequently, the pots were placed under greenhouse conditions to promote seed germination and were watered daily with the same amount of water until the emergence of one-leaf stage seedlings (Zadoks scale: GS11) [31]. At this stage, a thinning process was performed to leave one seedling in each pot, ensuring homogeneity with the remaining seedlings in the experiment. The pots were then irrigated with 1× Hoagland solution [32] to maintain the moisture content at “pot capacity” until the flag leaf emergence stage (Zadoks scale: GS37) [31], where the initiation of water deficit stress treatment started.

To assess the impact of water deficit stress on the four selected genotypes, water deficit treatments at two reproductive stages were imposed by allowing the pots to reach either 60% or 80% water depletion from their full “pot capacity” for intermediate and severe stress treatments, respectively. This was achieved by withholding the watering with 1× Hoagland solution and weighing the pots to achieve the targeted pot capacity for each stress

treatment. At these levels, the pots were kept without irrigation for three days before irrigating them again to reach a pot capacity of 60% and 40% for the intermediate and severe stress treatments, respectively. This cycle was repeated and maintained until the plants reached the anthesis growth stage (Zadoks scale: GS69) [31]. At that point, the water deficit stress regime was replaced with a “watering every 4 days” regime until physiological maturity, with 300 ml and 150 ml of water added for the intermediate and severe stress treatments, respectively. As a control treatment, the remaining pots were irrigated with 500 ml of 1× Hoagland solution to reach full pot capacity and were kept well-watered until the end of the experiment. For each treatment, the physiological responses of the treated plants were monitored three days after the onset of stress treatments at the two different growth stages: at the heading stage (Zadoks scale: GS59) and anthesis (Zadoks scale: GS69) [31]. Furthermore, agronomic measurements were taken at the end of the experiment, when the plants reached the physiological maturity stage (Zadoks scale: GS87) [31].

2.3. Agronomic Traits

Agronomic data were recorded to evaluate the performance of the three selected NILs and LND under water deficit stress. These measurements included: duration of heading (HD), calculated as the number of days from seed emergence to the point when the plant in the pot produced a fully developed spike (Zadoks scale: GS59) [31]; duration of physiological maturity (MD) was recorded as the number of days from seed emergence to the day when the plant in each plot had reached the physiological maturity stage (Zadoks scale: GS87) [31]; the grain-filling period (GFP) was determined by subtracting MD from HD; plant height (PH in cm) measured from the plant base to the top of the spike excluding awns; tiller number (TN) as the final number of tillers counted from each plant in the pot; spike number (SN) as the final number of spikes counted from each plant in the pot; spike length (SL in cm) measured from the spike base to the top excluding awns; awns length (AL in cm) measured from the apical spikelet to the top of awns; spikelet number (SpN) as the final number of spikelets per spike estimated as an average of three selected main spikes in the pot; fertile florets (FFL) as the number of fertile florets/spikelets estimated as an average of three selected main spikes in the pot; total weight (TW in g) measured as the total weight of the whole above-ground material including grains that was harvested from each pot; spikes weight (SW in g) measured in g by weighing the harvested spikes from each plant; grain weight (GW in g) measured by weighing the grains (after threshing) from each plant in the pot; grain number (GN) measured as the total number of grains counted from all harvested spikes from each plant in the pot; grains/spike (GpS) as the number of fertile spikelets/spike calculated as grain number divided on spike number; and thousand kernel weight (TGW in g) was analyzed using the MARVIN seed analyzer machine (Marvin-GTA, Sensorik GmbH, Version 5.0) using threshed grains from each plant in the pot.

2.4. Physiological Measurements

During the experiment, the physiological responses of the treated plants were assessed under water deficit and well-watered conditions. These physiological measurements were recorded at specific time points corresponding to the two different growth stages mentioned earlier. To determine the relative water content (RWC) of treated plants, a 5 cm piece was excised from a fully expanded leaf at the three selected growth stages. Immediately after excising the leaf, the fresh weight (FW) was recorded. Subsequently, the excised leaf piece was gently dried on a paper towel to remove excess water, and the turgid weight (TW) was recorded. Finally, the leaf samples were dried for 24 hours at 50 °C, and their dry weight (DW) was recorded. The RWC was calculated using the following formula, based on Barrs and Weatherley [33]: $RWC (\%) = [(FW - DW) / (TW - DW)] \times 100$.

Stomatal resistance ($s.m^{-1}$) was measured using a “steady-state” Porometer (AP4 model, Delta T devices, Cambridge, UK) attached to the abaxial side of the leaves at the two growth stages. Two readings per treatment were taken during midday (11:00 am-12:00 pm) on fully expanded leaves, and the average value was used for further analysis. Chlorophyll fluorescence (maximum yield of PSII (F_v/F_m)) was measured at the two growth stages with a pulse-modulated Fluorometer (OS1-FL modulated chlorophyll Fluorometer, ADC BioScientific Ltd., Hertford, UK). For relative measurement of chlorophyll content, a single-photon avalanche diode machine SPAD-250 (MC-100 Chlorophyll Meter, Apogee instruments, USA) was used to measure the chlorophyll content (SPAD index) using a fully expanded leaf with two readings, and the average value was used for further analysis.

2.5. Statistical Analysis

A split-plot design with four replications was used, where each replicate contained three subplots for well-watered, intermediate, and severe water deficit stress treatments, respectively. Each plot contained four genotypes randomly distributed in each subplot. Collected physiological and agronomical data were analyzed by using the GenStat program (Release 16.1, 2013; VSN International Ltd, UK). Analysis of variance (ANOVA) was performed for water deficit treatment and genotypes and their interactions, and the least significant difference test (LSD, $p \geq 0.05$) was used for mean separation. Pearson’s correlation coefficients were calculated in R using the corrplot package as described previously [34]. The pheatmap package v1.0.8 in R (<https://cran.r-project.org/web/packages/pheatmap/index>) was used to construct a heatmap and a hierarchical cluster using Euclidean distance and Ward.D2 method using the means of the tested genotypes.

3. RESULTS

3.1. Agronomic Data Analysis

The analysis of variance (ANOVA) indicated clear variation among water-deficit treatments for all agronomic traits, with the exception of SL (Table 1). Regarding the genotypic effect, significant variations were detected in 7 out of the 16 agronomic traits, specifically in MD, GFP, PH, TN, SN, FFL, and GpS. The interactive effect between genotype and treatment exhibited significant differences only in MD and GFP (Table 1). For the coefficient of variations (CV), the values varied for the agronomic traits where MD displayed the least variation, with a CV of 0.66%, and conversely, SN showed the highest variability with a CV of 14.43% (Table 1).

Table 1. Mean squares from combined analysis of variance for 16 agronomic traits in three durum wheat NILs and LND grown under three watering regimes.

Source of variation	DF	HD	MD	GFP	PH	SL	AL	TN	SN
Rep	3	0.02	0.03	0.02	1.83	1.01	1.02	0.41	0.53
Treatment	2	21.94 **	431.58**	264.15**	312.15**	5.14	35.51**	203.40**	175.69**
Error I	6	1.35	1.36	2.98	6.56	0.11	0.85	0.70	1.13
Genotype	3	0.58	189.69**	183.47**	28.94*	0.19	0.32	6.08**	7.14**
Treatment × Genotype	6	0.33	10.69**	8.51*	4.26	0.26	0.16	2.20	1.24
Error II	27	0.93	1.31	2.49	6.71	0.28	0.59	0.92	0.84
Grand Mean	-	140.06	177.04	36.98	127.83	9.27	9.81	7.77	7.38
C.V. (%)	-	0.83	0.66	4.67	2.00	3.52	9.41	10.78	14.43
Source of variation	DF	SpN	FFL	GpS	GN	TW	SW	GW	TGW
Rep	3	0.52	0.04	11.23	523.19	7.85	0.14	1.17	0.35
Treatment	2	16.01**	2.01**	757.14**	236189**	3816.38**	690.38**	387.01**	49.34**
Error I	6	0.93	0.07	7.08	513.82	5.09	1.42	1.84	0.57
Genotype	3	1.04	0.27**	124.32**	11.81	1.45	2.01	1.84	21.50
Treatment × Genotype	6	0.31	0.03	5.79	399.39	7.09	2.14	0.92	0.93
Error II	27	0.74	0.06	16.79	661.69	4.71	1.51	0.76	1.60
Grand Mean	-	22.87	2.36	31.78	235.07	30.82	12.94	9.44	39.71
C.V. (%)	-	4.21	11.28	8.37	9.64	7.32	9.21	14.36	1.91

Note: * is significant at $p \leq 0.05$ level; ** is significant at $p \leq 0.01$ level.




Fig. (1). Mean values of heading date (in green), maturity dates (in orange), and grain filling period (numbers within columns) for three tested durum wheat NILs and LND grown under three different watering regimes. Bars represent LSD test values at $p \leq 0.05$ to compare maturity date means ($n=4$) for the combined analysis and genotypes means across all treatments, while letters represent LSD test values at $p \leq 0.05$ for specific comparison of maturity date means ($n=4$) of the tested genotypes within each specific stress treatment.

No significant differences were detected for HD among the tested genotypes under various stress conditions (Fig. 1). Conversely, significant differences were evident for MD and GFP among the tested genotypes under different stress conditions. As expected, the NILs exhibited a significantly faster maturation rate with shorter GFP compared to LND (Fig. 1). Water stress treatments accelerated maturity in all tested genotypes, with a clear, pronounced effect on the tested NILs, especially as stress intensity increased, while no significant differences were observed for LND.

The deleterious effects of severe water deficit treatment were prominently evident in 13 agronomic traits, with a significant reduction in various parameters, including PH, SL, AL, TN, SN, SpN, FFL, GpS, GN, TW, SW, GW, and TGW, when compared to well-watered treatment conditions (Table 2). For instance, under severe water deficit treatment, the mean value of TW significantly decreased to 17.74 g compared to 26.87 g under moderate stress and 47.86 g in the well-watered treatment. A similar trend was observed in TGW, where it decreased to 37.97 g under severe water deficit treatment compared to 39.68 g under moderate stress and 41.49 g in the well-watered treatment. Spike number (SN) also exhibited a significant

decline, with a mean value of 11.19 spikes in the well-watered treatment compared to 5.19 spikes under severe water deficit conditions, showing no significant difference from the moderate water deficit treatment (5.75 spikes) (Table 2). The SW ranged from 19.96 g in the well-watered treatment to 6.94 g in the severe water deficit treatment, with significant differences between the three treatments. Similarly, GW varied from 14.59 g in the well-watered treatment to 4.80 g in the severe water deficit treatment. Additionally, GN demonstrated a substantial reduction with the well-watered treatment, yielding a mean value of 364.95, while only 124.19 grains were obtained under severe water deficit conditions (Table 2).

The genotypic effect analysis through ANOVA revealed significant differences in several agronomic traits, including PH, TN, SN, FFL, GpS, and TGW (Table 3). For instance, LND exhibited the highest mean values for both TGW (41.65 g) and GpS (36.56), which were significantly higher than the values observed in the three NILs carrying the *NAM-B1* gene (Table 3). On the other hand, LND produced significantly the lowest SN mean value compared to NILs carrying the *NAM-B1* gene. Furthermore, LND plants produced the lowest mean of PH at 125.67 cm when compared to the NIL plants (Table 3).

Table 2. Mean values of agronomic traits exhibited significant differences under the influence of water deficit treatment.

Treatment	HD	PH	SL	AL	TN	SN	SpN
Watered	141.37 a*	132.06 a	9.77 a	11.15 a	11.88 a	11.19 a	23.85 a
Moderate	139.68 b	128.19 b	9.38 a	10.08 a	6.00 b	5.75 b	22.90 b
Severe	139.12 b	123.25 c	8.65 b	8.21 b	5.44 b	5.19 b	21.85 c
Grand mean	140.06	127.83	9.27	9.81	7.77	7.38	22.87
^{LSD} (0.05)	1.01	2.22	0.28	0.80	0.72	0.92	0.83
Treatment	FFL	GpS	GN	TW	SW	GW	TGW
Watered	2.71 a	32.82 a	364.95 a	47.86 a	19.96 a	14.59 a	41.49 a
Moderate	2.37 b	38.08 b	216.08 b	26.87 b	11.91 b	8.95 b	39.68 b
Severe	2.00 c	24.44 c	124.19 c	17.74 c	6.94 c	4.80 c	37.97 c
Grand mean	2.36	31.78	235.07	30.82	12.94	9.44	39.71
^{LSD} (0.05)	0.23	2.30	19.61	1.95	1.03	1.17	0.66

Note: * Mean values (n=16) with different letters within the same column indicate statistically significant difference levels based on LSD test at $p \leq 0.05$.

Table 3. Mean values of agronomical traits with significant differences for the genotypic effect.

Genotype	PH	TN	SN	FFL	GpS	TGW
#504	129.00 a	8.33 a	7.83 a	2.28 b	30.24 b	39.42 b
IR17-47	128.92 a	7.83 a	7.50 a	2.28 b	30.74 b	38.57 b
IR51-8	127.75 ab	8.17 a	7.92 a	2.31 b	29.59 b	39.23 b
Langdon	125.67 b	6.75 b	6.25 b	2.58 a	36.56 a	41.65 a
Grand Mean	127.83	7.77	7.38	2.36	31.78	39.71
LSD (0.05)	2.17	0.80	0.77	0.21	3.43	1.06

Note: * Mean values (n=12) with different letters within the same column are significantly different at $p \leq 0.05$ level according to LSD test.

3.2. Physiological Data Analysis

The ANOVA revealed significant differences among water-deficit treatments for all physiological traits, except for SPAD at the heading stage, emphasizing the impact of water stress conditions on the measured parameters (Table 4). For the genotypic effect, significant variations were identified in five physiological traits that included SPAD and Fv/Fm , both assessed at the heading and anthesis stages, along with stomatal resistance (SR) at the anthesis stage. The interactive effect between genotype and treatment demonstrated significant differences only for SR at the anthesis stage (Table 4). The coefficient of variation (CV) values ranged from 3.31% to 22.14%, with the highest value

observed for SR at the heading stage, while the lowest was recorded for RWC at the heading stage.

Significant effects of water deficit treatment were observed on RWC and SR at both growth stages (Table 5), with stressed plants that displayed significant differences in their mean values compared to well-watered plants. For SPAD at the anthesis stage, no significant differences were observed between well-watered and moderate stress treatments, while the severe stress treatment produced significantly the highest mean value. The mean values of the maximum efficiency of photosystem II (Fv/Fm) at both growth stages were significantly higher in well-watered plants compared to those subjected to stress treatment.

Table 4. Mean squares from combined analysis of variance for eight physiological traits in three durum wheat NILs and LND grown under three watering regimes.

Source of variation	DF	RWC (H)	RWC (A)	SPAD (H)	SPAD (A)	Fv/Fm (H)	Fv/Fm (A)	SR (H)	SR (A)
Rep	3	1.43	8.58	194.66	549.83	5.49 ^{e-04}	2.11 ^{e-04}	2.54	2.74
Treatment	2	799.23**	1429.45**	1246.27	5102.21**	19.2 ^{e-03**}	0.023**	985.21**	4431.84**
Error I	6	8.03	27.02	545.00	446.50	2.14^{e-04}	9.13^{e-04}	3.15	4.24
Genotype	3	10.55	2.70	1693.21*	8918.62**	1.48 ^{e-03**}	2.61 ^{e-04**}	1.86	66.24**
Treatment × Genotype	6	2.76	5.78	80.95	217.18	4.26 ^{e-04}	1.31 ^{e-03}	1.46	66.44**
Error II	27	9.82	7.98	378.25	379.97	2.95^{e-04}	5.06^{e-04}	1.89	6.10
Grand Mean		85.55	6.39	422.68	403.07	0.59	0.60	8.02	15.05
C.V. (%)		3.31	5.20	5.52	5.24	2.46	5.05	22.14	13.69

Note: * is significant at $p \leq 0.05$ level; ** is significant at $p \leq 0.01$ level. H: Heading growth stage; A: Anthesis growth stage.

Regarding the genotypic effect, SPAD mean values for NIL #504 were significantly higher than those of other tested genotypes at both growth stages (Table 6). For *Fv/Fm*, LND consistently produced the highest mean values at both growth stages compared to NILs carrying

the *NAM-B1* gene. On the other hand, LND produced the lowest mean value of SR at the anthesis stage, which was statistically significant when compared with the other tested genotypes (Table 6).

Table 5. Mean values of physiological traits exhibited significant differences under the influence of water deficit treatment.

Treatment	RWC (H)	RWC (A)	SPAD (A)	<i>Fv/Fm</i> (H)	<i>Fv/Fm</i> (A)	SR (H)	SR (A)
Watered	93.53 a*	91.38 a	387.23 a	0.63 a	0.64 a	1.43 a	2.29 a
Moderate	83.02 b	80.01 b	399.56 a	0.59 b	0.58 b	5.93 b	8.98 b
Severe	80.09 c	72.62 c	422.42 b	0.56 b	0.57 b	16.70 c	33.88 c
Grand mean	85.55	81.34	403.07	0.59	0.60	8.02	15.05
LSD(0.05)	2.45	4.5	18.28	0.01	0.02	1.54	1.78

Note: * Mean values (n=16) with different letters within the same column are significantly different at $p \leq 0.05$ level according to LSD test. H: Heading growth stage; A: Anthesis growth stage.

Table 6. Mean values of physiological traits with significant differences for genotypic effect.

Genotype	SPAD (H)	SPAD (A)	<i>Fv/Fm</i> (H)	<i>Fv/Fm</i> (A)	SR (A)
#504	440.38 a*	443.71 a	0.59 b	0.60 b	15.91 a
IR17-47	415.93 b	393.72 b	0.59 b	0.59 b	17.19 a
IR51-8	415.68 b	388.21 b	0.59 b	0.59 b	15.38 a
Langdon	418.74 b	386.64 b	0.61 a	0.62 a	11.72 b
Overall Mean	422.68	403.07	0.59	0.60	15.05
LSD (0.05)	16.29	16.33	0.01	0.03	2.07

Note: * Mean values (n=12) with different letters within the same column are significantly different at $p \leq 0.05$ level according to LSD test. H: Heading growth stage; A: Anthesis growth stage.




Fig. (2). Mean values of stomata resistance at the anthesis stage for three tested durum wheat NILs and LND grown under three different watering regimes. Bars represent LSD test values at $p \leq 0.05$ to compare stomata resistance means for the combined analysis and genotypes means (n=4) across all treatments, while letters represent LSD test values at $p \leq 0.05$ for specific comparison of stomata resistance means (n=4) of the tested genotypes within each specific stress treatment.

Concerning the genotype × treatment interactive effect, only SR at the anthesis stage demonstrated statistical significance, with the mean values of the tested genotypes increasing with the severity of the stress treatment (Fig. 2). Notably, under severe stress conditions, NILs carrying *NAM-B1* exhibited significantly higher mean values than LND, indicating a lower transpiration rate in the NILs.

3.3. Correlation and Heatmap Analysis

In general, significant positive correlations ($p \geq 0.05$) were detected among the examined traits, with negative correlations primarily observed between the tested traits and SR (Fig. 3a). The MD exhibited a positive correlation

with all parameters except PH, TN, SN, and SPAD at the heading stage. Furthermore, GW showed positive correlations with all traits, excluding G/S, SPAD, and SR. The heatmap cluster analysis showed three 303 distinct clusters aligned with the stress treatment levels, remarkably out-grouping LND within each cluster (Fig. 3b). From this perspective, higher estimated values for LND were obvious for MD, GFP, FFL, and TGW regardless of the stress treatment. Additionally, a clear increase in *Fv/Fm* values was observed only under severe stress treatment. Interestingly, NIL #504 exhibited higher estimated values for SPAD, showing a substantial increase with rising stress levels (Fig. 3b).




Fig. 5 contd.....




Fig. (3). (a) Pearson's coefficients and pairwise correlations among traits in three tested durum wheat NILs and LND under three distinct watering regimes (blank squares denote non-significance at $p \leq 0.05$ level); (b) Heatmap clustering based on the data from three tested durum wheat NILs and LND cultivated under three distinct watering regimes (C: well-watered; MS: moderate stress; SS: severe stress). The color intensity in each figure reflects the magnitude of the corresponding estimate.

4. DISCUSSION

In this study, we investigated the role of *NAM-B1* in durum wheat in response to water deficit stress conditions at two reproductive stages. For this purpose, a selected set of durum wheat (*Triticum turgidum* spp. *durum*) NILs

differing in the *NAM-B1* gene was used. The check genotype (LND) was the only genotype that did not carry the *NAM-B1* gene, while the other three lines carried chromosomal segments from *Triticum turgidum* ssp. *dicoccoides* with the *NAM-B1* functional allele. *NAM-B1* is a well-studied gene in wheat that encodes a transcription

factor involved in accelerating senescence and enhancing protein, iron, and zinc content in grains [35]. Recently, it was found to act as a positive regulator of nutrient translocation from vegetative tissues to grains [36]. *NAM-B1* gene plays diverse roles, including nutrition remobilization, enhancing grain protein content, altering grain micronutrient content, and accelerating monocarpic senescence in wheat [20, 37]. The *NAM-B1* expression was detected predominantly in the flag leaf, declining at early booting and post-anthesis stages [38].

The study revealed significant differences in several agronomical and physiological traits in response to stress treatment, indicating clear negative effects of water deficit across genotypes (Table 1). Water deficit stress led to reductions in multiple traits, including GW, GN, TW, PH, and TN, which is consistent with previous findings [39]. These detrimental effects on wheat agronomical and physiological performance, particularly yield, are well-established [40]. Similar negative effects under stress conditions were observed in previous studies for various agronomic traits [41]. Leaf water potential, osmotic potential, and relative water content also exhibited significant reductions under drought stress [42]. Physiological traits like transpiration rate, chlorophyll content, and RWC were significantly reduced under post-anthesis drought stress in wheat [43]. In contrast, genotypic effects were less pronounced in this study, with few agronomical and physiological traits affected. Inconsistent responses to water deficit were observed in previous studies, highlighting the complexity of drought tolerance traits influenced by various genes and environmental interactions [40, 44].

The PH of genotypes varied significantly among water deficit treatments, with the genotypes exhibiting a reduction under severe water deficit, consistent with previous findings on the pronounced effect of drought stress on PH in durum wheat [39]. On the other hand, NILs carrying *NAM-B1* showed variable responses, where #504 and IR17-47 showed significant differences when compared with LND, while IR51-8 showed no significant differences. In a previous study, the derivatives lines resulting from the introgression of the *NAM-B1* gene in the wheat cultivar "HUU468" showed variable plant height mean values that ranged from 80-92 cm when compared to 85.7 cm of the parental line [45], which is in general agreement with our results. For SL, LND produced the lowest mean value under severe water deficit treatment, although not significantly different, compared to NILs carrying *NAM-B1*. The agronomic performance of BC2F3 families carrying the *NAM-B1* gene identified a single line (HUU468-09-6) from several families with significantly higher spike length mean values when compared to the recurrent parent HUU468 [45].

For TN and SN, a clear significant reduction under water deficit conditions was observed, with LND showing significantly lower mean values than NILs carrying *NAM-B1*. The SN per plant in lines carrying the functional *NAM-B1* allele exhibited an average increase of 4.5 spikes per plant when compared with LND plants [45, 46]. Kumar *et al.* [28] showed that genotypes carrying *NAM-B1* did not differ significantly in TN, while the introgression of the functional allele of *NAM-B1* increased SN when compared with the check cultivar under field conditions [20]. It is well established that the *NAM-B1* gene is highly expressed only after the anthesis stage [35]. Therefore, it is unexpected to see an effect of the *NAM-B1* gene on SN and TN, as these traits are usually determined before anthesis [47]. However, the positive impact of the *NAM-B1* gene on SN and TN and other traits needs further investigation to confirm the allelic effect in introgressed lines.

The total weight of all tested genotypes was affected by water deficit treatment, with no clear significant differences between the tested genotypes (Table 2). Drought is known to decrease dry weight in durum wheat lines [48]. Deckard *et al.* [49] studied the responses of Langdon [LDN(DIC)] (carrying *NAM-B1*) to N-uptake under field conditions and found that higher accumulation ratios (N-uptake to TW) were a result of a lower TW in substitution lines when compared with LND, which was not observed under greenhouse conditions in this study. The GW and GN showed no significant differences between genotypes under different treatments but were reduced under water deficit conditions. Positive correlations observed between GW and several traits align well with earlier studies [50]. For GN, Kaznina *et al.* [24] studied the effect of Zn deficiency on GN and GW in wheat lines carrying different allelic variants of the *NAM-B1* gene, and they found a significant increase in GN and GW only under Zn stress when compared to the control. Under water deficit stress, senescence can be accelerated, leading to increased nutrient remobilization and higher GPC and micronutrient concentrations in grains [37].

Grain per spike results indicated that LND produced the highest mean values under well-watered and severe water deficit conditions (Table 2). For TGW, a significant reduction in NILs carrying *NAM-B1* was observed, consistent with previous reports [28, 51]. On the other hand, Eagles *et al.* [51] showed that NIL-carrying *NAM-B1* had a negligible effect on grain yield when compared with their recurrent parents in field trials under Australian environments. They suggested crossing newly commercial cultivars, which are introgressed with *NAM-B1*, with large grain size genotypes with targeted selection for grain weight.

Physiological parameters were negatively affected by water deficit treatment. For instance, SR significantly increased in response to water deficit conditions, consistent with previous findings [52]. Chlorophyll fluorescence (*Fv/Fm*) differences between water treatments at different growth stages reflect the complex response of the tested genotypes [53]. Water deficits led to a significant increase in SR, supporting the notion that genotypes carrying the *NAM-B1* gene play a role in accelerating senescence, particularly under severe water deficits [25]. At the anthesis stage, NIL #504 showed significantly higher mean values of SPAD when compared with other tested genotypes. Akhka *et al.* [54] studied the

effect of water deficit on chlorophyll content in tested durum wheat cultivars, and they observed a reduction in chlorophyll content in all tested cultivars under drought conditions. A recurrent study of BC2 NILs carrying a major QTL for TGW on 6A for two growing seasons showed no significant differences in relative chlorophyll content after 20–25 days of anthesis, and significant differences were detected after 45 days of anthesis [55]. Botyanszka *et al.* [53] reported that the mean values of chlorophyll contents in drought-tolerance genotypes of barley were significantly higher than those in drought-sensitive genotypes under water deficit. On the other hand, genotypes that carry the functional *NAM-B1* gene are known to play a major role in accelerating senescence [23], which was observed in this study and, in particular, in response to severe water deficit treatment. This could be attributed to the growth stages selected in this study, where water deficit was imposed and continued till the later stages of wheat development.

In this study, the majority of the studied traits were not affected by the treatment × genotype interactions, and this might be related to the complexity of the drought tolerance trait, which is described as a quantitative trait controlled by many genes with their expression influenced by different environmental and genetic interactions [56]. The lack of significance for the interactive effect might be explained by the masked impact of a single QTL selected in this study in the NIL background. Furthermore, this may highlight the necessity for further investigations into the role of *NAM-B1* under different growth stages and environmental conditions. Another possibility might be related to the inducible expression of *NAM-B1* in response to drought or to the growth stage. The expression of *NAM-A1*, the *NAM-B1* closest homolog in wheat, was found to be negatively affected by heat at senescence, while no significant effect of drought stress on *NAM-A1* expression was observed [30]. This is also supported by the findings of Kaul *et al.* [57], who found that the expression of *NAM-B1* was negligibly at the seedling and vegetative stages and was highly expressed at the senescence stage. Recently, a stress-inducible allele of *NAM-A1* (*TaSNAC8-6A*) was identified, which was induced in response to stresses at the seedling stage [29]. A three-bp indel in the promoter region resulted in a stress-responsive *cis*-regulatory element in the *NAM-A1* promoter that induced drought tolerance in selected genotypes. In this study, the size of introgression in selected NILs varied, and this could explain the observed variations among them in terms of different traits. Future studies are needed to fine-map minor QTLs associated with responses against drought in tested NILs or to dissect the role of *NAM-B1* under stress at different growth stages.

CONCLUSION

This study investigated the combined effects of genotype and water deficit stress in two reproductive stages on agronomic and physiological traits in three NILs carrying the *NAM-B1* gene compared to their recurrent

parent (LND) in durum wheat. Our findings revealed significant variations in both agronomic and physiological traits in response to water stress treatment or genotype. The genotypic effect was pronounced in seven agronomic traits, including MD, GFP, PH, TN, SN, FFL, and GpS, with NILs carrying the *NAM-B1* gene exhibiting, as expected, a faster maturation rate and shorter GFP compared to LDN. Severe water deficit significantly reduced various agronomic parameters, such as TW, SW, GW, and TGW, emphasizing the detrimental effects of severe stress conditions on reproductive stages in durum wheat plants. Further, physiological traits, in particular SR and *Fv/Fm*, displayed significant variations among genotypes and stress treatments, with NILs exhibiting clear differences for SR at the anthesis stage under severe stress conditions compared to LND. Nonetheless, this study highlights the need for further research to elucidate the underlying genetic mechanisms mediated by *NAM* genes in response to stress conditions across different growth stages.

LIST OF ABBREVIATIONS

RWC	=	Relative Water Content
DW	=	Dry Weight
TW	=	Turgid Weight
FW	=	Fresh Weight
ROS	=	Reactive Oxygen Species

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

The datasets supporting the results of this article will be freely available upon reasonable request from the corresponding author [A.A-A].

FUNDING

This study was funded by Deanship of Scientific Research, The University of Jordan.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Food and Agriculture Organization statistics [FAOSTAT]. FAO Stat 2021; FAO, Rome. 2021. Available from: <http://www.fao.org/faostat>
- [2] Shewry PR, Hey SJ. The contribution of wheat to human diet and health. *Food Energy Secur* 2015; 4(3): 178-202. <http://dx.doi.org/10.1002/fes3.64> PMID: 27610232
- [3] International Grains Council [IGC]. 2016; Online Database: World Grain Statistics. 2016. Available from: <https://www.igc.int/en/subscriptions/subscription> (accessed on 21 May 2023).
- [4] Fu BX, Wang K, Dupuis B, Taylor D, Nam S. Kernel vitreousness and protein content: Relationship, interaction and synergistic effects on durum wheat quality. *J Cereal Sci* 2018; 79: 210-7.

- <http://dx.doi.org/10.1016/j.jcs.2017.09.003>
- [5] Saini P, Kaur H, Tyagi V, et al. Nutritional value and end-use quality of durum wheat. *Cereal Res Commun* 2023; 51(2): 283-94. <http://dx.doi.org/10.1007/s42976-022-00305-x>
- [6] Ozturk M, Gul A. Climate change and food security with emphasis on wheat. Academic Press 2020; pp. 43-51.
- [7] Lobell DB, Sibley A, Ivan Ortiz-Monasterio J. Extreme heat effects on wheat senescence in India. *Nat Clim Chang* 2012; 2(3): 186-9. <http://dx.doi.org/10.1038/nclimate1356>
- [8] Yadav AK, Carroll AJ, Estavillo GM, Rebetzke GJ, Pogson BJ. Wheat drought tolerance in the field is predicted by amino acid responses to glasshouse-imposed drought. *J Exp Bot* 2019; 70(18): 4931-48. <http://dx.doi.org/10.1093/jxb/erz224> PMID: 31189018
- [9] Farooq M, Hussain M, Siddique KHM. Drought stress in wheat during flowering and grain-filling periods. *Crit Rev Plant Sci* 2014; 33(4): 331-49. <http://dx.doi.org/10.1080/07352689.2014.875291>
- [10] Prasad PVV, Pisipati SR, Momčilović I, Ristic Z. Independent and combined effects of high temperature and drought stress during grain filling on plant yield and chloroplast EF-Tu expression in spring wheat. *J Agron Crop Sci* 2011; 197(6): 430-41. <http://dx.doi.org/10.1111/j.1439-037X.2011.00477.x>
- [11] Dhanda SS, Sethi GS. Tolerance to drought stress among selected Indian wheat cultivars. *J Agric Sci* 2002; 139(3): 319-26. <http://dx.doi.org/10.1017/S0021859602002526>
- [12] Anjum MM, Arif M, Riaz M, Akhtar K, Zhang SQ, Zhao CP. Performance of hybrid wheat cultivars facing deficit irrigation under semi-arid climate in Pak. *Agronomy* 2021; 11(10): 1976. <http://dx.doi.org/10.3390/agronomy11101976>
- [13] Kulkarni M, Soolanayakanahally R, Ogawa S, Uga Y, Selvaraj MG, Kagale S. Drought response in wheat: key genes and regulatory mechanisms controlling root system architecture and transpiration efficiency. *Front Chem* 2017; 5: 106. <http://dx.doi.org/10.3389/fchem.2017.00106> PMID: 29259968
- [14] Manna M, Thakur T, Chirom O, Mandlik R, Deshmukh R, Salvi P. Transcription factors as key molecular target to strengthen the drought stress tolerance in plants. *Physiol Plant* 2021; 172(2): 847-68. <http://dx.doi.org/10.1111/pp1.13268> PMID: 33180329
- [15] Singh S, Koyama H, Bhati KK, Alok A. The biotechnological importance of the plant-specific NAC transcription factor family in crop improvement. *J Plant Res* 2021; 134(3): 475-95. <http://dx.doi.org/10.1007/s10265-021-01270-y> PMID: 33616799
- [16] Tang Y, Liu M, Gao S, et al. Molecular characterization of novel TaNAC genes in wheat and overexpression of TaNAC2a confers drought tolerance in tobacco. *Physiol Plant* 2012; 144(3): 210-24. <http://dx.doi.org/10.1111/j.1399-3054.2011.01539.x> PMID: 22082019
- [17] Zhang Z, Liu C, Guo Y. Wheat transcription factor TaSNAC11-4B positively regulates leaf senescence through promoting ROS production in transgenic Arabidopsis. *Int J Mol Sci* 2020; 21(20): 7672. <http://dx.doi.org/10.3390/ijms21207672> PMID: 33081330
- [18] Hu XG, Wu BH, Liu DC, Wei YM, Gao SB, Zheng YL. Variation and their relationship of NAM-G1 gene and grain protein content in *Triticum timopheevii* Zhuk. *J Plant Physiol* 2013; 170(3): 330-7. <http://dx.doi.org/10.1016/j.jplph.2012.10.009> PMID: 23218544
- [19] Al Abdallat AM, Ayad JY, Abu Elenein JM, Al Ajlouni Z, Harwood WA. Overexpression of the transcription factor HvSNAC1 improves drought tolerance in barley (*Hordeum vulgare* L.). *Mol Breed* 2014; 33(2): 401-14. <http://dx.doi.org/10.1007/s11032-013-9958-1>
- [20] Ma J, Tang X, Sun B, et al. A NAC transcription factor, TaNAC5D-2, acts as a positive regulator of drought tolerance through regulating water loss in wheat (*Triticum aestivum* L.). *Environ Exp Bot* 2022; 196: 104805. <http://dx.doi.org/10.1016/j.envexpbot.2022.104805>
- [21] Mao H, Li S, Chen B, et al. Variation in cis-regulation of a NAC transcription factor contributes to drought tolerance in wheat. *Mol Plant* 2022; 15(2): 276-92. <http://dx.doi.org/10.1016/j.molp.2021.11.007> PMID: 34793983
- [22] Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J. A NAC Gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science* 2006; 314(5803): 1298-301. <http://dx.doi.org/10.1126/science.1133649> PMID: 17124321
- [23] Tabbita F, Pearce S, Barneix AJ. Breeding for increased grain protein and micronutrient content in wheat: Ten years of the GPC-B1 gene. *J Cereal Sci* 2017; 73: 183-91. <http://dx.doi.org/10.1016/j.jcs.2017.01.003>
- [24] Kaznina N, Dubovets N, Batova Y, Ignatenko A, Orlovskaya O, Repkina N. The response of wheat with different allele statuses of the Gpc-B1 gene under zinc deficiency. *Agronomy* 2021; 11(6): 1057. <http://dx.doi.org/10.3390/agronomy11061057>
- [25] Lundström M, Leino MW, Hagenblad J. Evolutionary history of the NAM-B1 gene in wild and domesticated tetraploid wheat. *BMC Genet* 2017; 18(1): 118. <http://dx.doi.org/10.1186/s12863-017-0566-7> PMID: 29262777
- [26] Hale I, Zhang X, Fu D, Dubcovsky J. Registration of wheat lines carrying the partial stripe rust resistance gene Yr36 without the Gpc-B1 high grain protein content allele. *J Plant Regist* 2013; 7(1): 108-12. <http://dx.doi.org/10.3198/jpr2012.03.0150crg> PMID: 26962384
- [27] Velu G, Singh RP, Cardenas ME, Wu B, Guzman C, Ortiz-Monasterio I. Characterization of grain protein content gene (GPC-B1) introgression lines and its potential use in breeding for enhanced grain zinc and iron concentration in spring wheat. *Acta Physiol Plant* 2017; 39(9): 212. <http://dx.doi.org/10.1007/s11738-017-2509-3>
- [28] Kumar J, Jaiswal V, Kumar A, et al. Introgression of a major gene for high grain protein content in some Indian bread wheat cultivars. *Field Crops Res* 2011; 123(3): 226-33. <http://dx.doi.org/10.1016/j.fcr.2011.05.013>
- [29] Mao H, Li S, Wang Z, et al. Regulatory changes in TaSNAC8-6A are associated with drought tolerance in wheat seedlings. *Plant Biotechnol J* 2020; 18(4): 1078-92. <http://dx.doi.org/10.1111/pbi.13277> PMID: 31617659
- [30] Al-Sayaydeh R, Al-Hababeh K, Akkeh Z, Albdaiwi RN. *In silico* gene expression analysis of the stress-related NAC-A gene subfamily to dissect their role in abiotic stress tolerance in bread wheat (*Triticum aestivum* L.). *J Agric Sci* 2021; 17(3): 341-54. <http://dx.doi.org/10.35516/ijas.v17i3.90>
- [31] Zadoks JC, Chang TT, Konzak CF. A decimal code for the growth stages of cereals. *Weed Res* 1974; 14(6): 415-21. <http://dx.doi.org/10.1111/j.1365-3180.1974.tb01084.x>
- [32] Hoagland DR, Arnon DI. The water-culture method for growing plants without soil. *Circular* 1950; 347: 32.
- [33] Barrs HD, Weatherley PE. A re-examination of the relative turgidity technique for estimating water deficit in leaves. *Aust J Biol Sci* 1962; 15(3): 413-28. <http://dx.doi.org/10.1071/BI9620413>
- [34] Al-Sayaydeh R, Shtaya MJ, Qubbaj T, et al. Performance and stability analysis of selected durum wheat genotypes differing in their kernel characteristics. *Plants* 2023; 12(14): 2664. <http://dx.doi.org/10.3390/plants12142664> PMID: 37514278
- [35] Uauy C, Brevis JC, Dubcovsky J. The high grain protein content gene Gpc-B1 accelerates senescence and has pleiotropic effects on protein content in wheat. *J Exp Bot* 2006; 57(11): 2785-94. <http://dx.doi.org/10.1093/jxb/erl047> PMID: 16831844
- [36] Alhabbar Z, Islam S, Yang R, et al. Associations of NAM-A1 alleles with the onset of senescence and nitrogen use efficiency under Western Australian conditions. *Euphytica* 2018; 214(10): 180. <http://dx.doi.org/10.1007/s10681-018-2266-4>
- [37] Shoormij F, Mirlohi A, Chan-Rodriguez D, Bolibok-Braęoszewska H, Saeidi G. Characterization of 14 *Triticum* species for the NAM-B1 gene and its associated traits. *PLoS One* 2023; 18(8): e0287798. <http://dx.doi.org/10.1371/journal.pone.0287798> PMID: 37607184
- [38] Bhattacharjee S. Molecular cloning and gene expression profiling

- of GPC-B1 (Grain Protein Content B1) under terminal heat stress in wheat (*Triticum aestivum* L.). PhD Thesis, Division of Molecular Biology and Biotechnology Icar-Indian Agricultural Research Institute 2017.
- [39] Liu H, Searle IR, Mather DE, Able AJ, Able JA. Morphological, physiological and yield responses of durum wheat to pre-anthesis water-deficit stress are genotype-dependent. *Crop Pasture Sci* 2015; 66(10): 1024-38.
<http://dx.doi.org/10.1071/CP15013>
- [40] Gupta NK, Gupta S, Kumar A. Effect of water stress on physiological attributes and their relationship with growth and yield of wheat cultivars at different stages. *J Agron Crop Sci* 2001; 186(1): 55-62.
<http://dx.doi.org/10.1046/j.1439-037x.2001.00457.x>
- [41] Mostafa A, Ali N, Hossein S, Mehdi H. Effects of drought stress on some agronomic and morphological traits of durum wheat (*Triticum durum* Desf.) landraces under greenhouse condition. *Afr J Biotechnol* 2011; 10(64): 14097-107.
<http://dx.doi.org/10.5897/AJB11.2322>
- [42] Tshikunde NM, Odindo A, Shimelis H, Mashilo J. Leaf gas exchange and water-use efficiency of dry-land wheat genotypes under water stressed and non-stressed conditions. *Acta Agric Scand - B Soil Plant Sci* 2018; 68(8): 738-48.
<http://dx.doi.org/10.1080/09064710.2018.1480729>
- [43] Akman H, Zhang C, Ejeta G. Physio-morphological, biochemical, and anatomical traits of drought-tolerant and susceptible sorghum cultivars under pre- and post-anthesis drought. *Physiol Plant* 2021; 172(2): 912-21.
<http://dx.doi.org/10.1111/ppl.13242> PMID: 33063861
- [44] Al-Ajlouni Z, Al-Abdallat A, Al-Ghzawi A, *et al.* Impact of pre-anthesis water deficit on yield and yield components in barley (*Hordeum vulgare* L.) plants grown under controlled conditions. *Agronomy* 2016; 6(2): 33.
<http://dx.doi.org/10.3390/agronomy6020033>
- [45] Vishwakarma MK, Mishra VK, Gupta PK, Yadav PS, Kumar H, Joshi AK. Introgression of the high grain protein gene *Gpc-B1* in an elite wheat variety of Indo-Gangetic Plains through marker assisted backcross breeding. *Curr Plant Biol* 2014; 1: 60-7.
<http://dx.doi.org/10.1016/j.cpb.2014.09.003>
- [46] Vishwakarma MK, Arun B, Mishra VK, Yadav PS, Kumar H, Joshi AK. Marker-assisted improvement of grain protein content and grain weight in Indian bread wheat. *Euphytica* 2016; 208(2): 313-21.
<http://dx.doi.org/10.1007/s10681-015-1598-6>
- [47] González FG, Miralles DJ, Slafer GA. Wheat floret survival as related to pre-anthesis spike growth. *J Exp Bot* 2011; 62(14): 4889-901.
<http://dx.doi.org/10.1093/jxb/err182> PMID: 21705386
- [48] Boutraa T, Akhkha A, Al-Shoabi AA, Alhejeli AM. Effect of water stress on growth and water use efficiency (WUE) of some wheat cultivars (*Triticum durum*) grown in Saudi Arabia. *J Taibah Univ Sci* 2010; 3(1): 39-48.
[http://dx.doi.org/10.1016/S1658-3655\(12\)60019-3](http://dx.doi.org/10.1016/S1658-3655(12)60019-3)
- [49] Deckard EL, Joppa LR, Hammond JJ, Hareland GA. Grain protein determinants of the Langdon durum-*dicoccoides* chromosome substitution lines. *Crop Sci* 1996; 36(6): 1513-6.
<http://dx.doi.org/10.2135/cropsci1996.0011183X003600060017x>
- [50] Brevis JC, Chicaiza O, Khan IA, Jackson L, Morris CF, Dubcovsky J. Agronomic and quality evaluation of common wheat near-isogenic lines carrying the leaf rust resistance gene *Lr47*. *Crop Sci* 2008; 48(4): 1441-51.
<http://dx.doi.org/10.2135/cropsci2007.09.0537>
- [51] Eagles HA, McLean R, Eastwood RF, *et al.* High-yielding lines of wheat carrying Gpc-B1 adapted to Mediterranean-type environments of the south and west of Australia. *Crop Pasture Sci* 2014; 65(9): 854-61.
<http://dx.doi.org/10.1071/CP14106>
- [52] El Hafid R, Smith DH, Karrou M, Samir K. Physiological responses of spring durum wheat cultivars to early-season drought in a Mediterranean environment. *Ann Bot* 1998; 81(2): 363-70.
<http://dx.doi.org/10.1006/anbo.1997.0567>
- [53] Botyanszka L, Zivcak M, Chovancek E, *et al.* Chlorophyll fluorescence kinetics may be useful to identify early drought and irrigation effects on photosynthetic apparatus in field-grown wheat. *Agronomy* 2020; 10(9): 1275.
<http://dx.doi.org/10.3390/agronomy10091275>
- [54] Akhkha A, Boutraa T, Alhejely A. The rates of photosynthesis, chlorophyll content, dark respiration, proline and abscisic acid (ABA) in wheat (*Triticum durum*) under water deficit conditions. *Int J Agric Biol* 2011; 13(2): 215-21.
[http://dx.doi.org/10.1016/S1658-3655\(12\)60019-3](http://dx.doi.org/10.1016/S1658-3655(12)60019-3)
- [55] Simmonds J, Scott P, Leverington-Waite M, *et al.* Identification and independent validation of a stable yield and thousand grain weight QTL on chromosome 6A of hexaploid wheat (*Triticum aestivum* L.). *BMC Plant Bio* 2014; 14(1)
<http://dx.doi.org/10.1186/s12870-014-0191-9>
- [56] Fleury D, Jefferies S, Kuchel H, Langridge P. Genetic and genomic tools to improve drought tolerance in wheat. *J Exp Bot* 2010; 61(12): 3211-22.
<http://dx.doi.org/10.1093/jxb/erq152> PMID: 20525798
- [57] Kaul T, Eswaran M, Ahmad S, *et al.* Probing the effect of a plus 1bp frameshift mutation in protein-DNA interface of domestication gene, *NAMBI*, in wheat. *J Biomol Struct Dyn* 2020; 38(12): 3633-47.
<http://dx.doi.org/10.1080/07391102.2019.1680435> PMID: 31621500